

**SURVIVAL, GROWTH, AND RADULA MORPHOLOGY OF POSTLARVAL PINTO
ABALONE (*HALIOTIS KAMTSCHATKANA*) WHEN FED SIX SPECIES OF BENTHIC
DIATOMS**

By

Lillian Miller Kuehl

Accepted in Partial Completion of
the Requirements for the Degree
Master of Science

ADVISORY COMMITTEE

Dr. Deborah Donovan, Chair

Dr. Brian Bingham

Dr. Suzanne Strom

GRADUATE SCHOOL

David L. Patrick, Interim Dean

MASTER'S THESIS

In presenting this thesis in partial fulfillment of the requirements for a master's degree at Western Washington University, I grant to Western Washington University the non-exclusive royalty-free right to archive, reproduce, distribute, and display the thesis in any and all forms, including electronic format, via any digital library mechanisms maintained by WWU.

I represent and warrant this is my original work, and does not infringe or violate any rights of others. I warrant that I have obtained written permissions from the owner of any third party copyrighted material included in these files.

I acknowledge that I retain ownership rights to the copyright of this work, including but not limited to the right to use all or part of this work in future works, such as articles or books.

Library users are granted permission for individual, research, and non-commercial reproduction of this work for educational purposes only. Any further digital posting of this document requires specific permission from the author.

Any copying or publication of this thesis for commercial purposes, or for financial gain, is not allowed without my written permission.

Lillian Miller Kuehl

April 22, 2020

**SURVIVAL, GROWTH, AND RADULA MORPHOLOGY OF POSTLARVAL PINTO
ABALONE (*HALIOTIS KAMTSCHATKANA*) WHEN FED SIX SPECIES OF BENTHIC
DIATOMS**

A Thesis
Presented to
The Faculty of
Western Washington University

In Partial Fulfillment
Of the Requirements for the Degree
Master of Science

by
Lillian Miller Kuehl
April 22, 2020

ABSTRACT

Haliotis kamtschatkana Jonas (pinto or northern abalone) is the only abalone native to the Pacific Northwest of North America. *Haliotis kamtschatkana* populations are in decline, and current restoration efforts in Washington State rely on out-planting hatchery-produced juveniles. Although several other abalone species are cultured extensively, little information exists on the cultivation of *H. kamtschatkana*, and hatchery production of this species has largely been a matter of trial and error. Hatcheries report highest mortalities in the postlarval stage, especially the first 3 to 6 months. Postlarvae feed on films of benthic diatoms, and the purpose of this study was to test 6 benthic diatom species as suitable diatom diets for *H. kamtschatkana*. Diatom diet suitability might rely on several factors, including morphology of the radula. The radula is a crucial feeding structure for gastropods and may display morphological plasticity, but it has never been characterized in *H. kamtschatkana* postlarvae. We investigated survival, growth, and radula morphology of *H. kamtschatkana* postlarvae when fed one of 6 benthic diatom species for 61 days post-settlement. *Amphora salina* best supported survival, especially in the first 20 days post-settlement (mean of 60% [SD, 22%] at day 20, mean of 47% [SD, 16%] at day 61), and *Achnanthes brevipes* yielded exceptionally low survival (mean of 12% [SD, 13%] and day 20, mean of 1% [SD, 3%] at day 61). Postlarvae fed *Cylindrotheca closterium* grew fastest among treatments (linear mixed model shell length = $293 * e^{0.021t}$, measured 1,110 μm [SD, 244 μm] at day 61), followed by postlarvae fed *Amphora salina*, *Navicula incerta*, or *Nitzschia laevis* (no significant difference between these diets; linear mixed model shell length = $302 * e^{0.018t}$, measured 894 μm [SD, 132 μm] at day 61).

We found no effect of diatom diet on radula morphology, but morphology was similar to that of other abalone species, with similar correlations between morphological characteristics and

shell length. We recommend that radula development of other species may be used as a proxy for *H. kamtschatkana* radula development, in the absence of further investigation.

We recommend *A. salina* as a suitable diet for newly settled *H. kamtschatkana* postlarvae, and that a combination of *A. salina* and *C. closterium* be investigated to support both survival and growth.

ACKNOWLEDGEMENTS

I would like to thank Puget Sound Restoration Fund for their donation of abalone and algae to this project, and particularly Josh Bouma for invaluable abalone advice and Stuart Ryan for algal culture. I received generous funding from the Pacific Northwest Shell Club Malacological Scholarship, the National Shellfisheries Association Michael Castagna Student Grant for Applied Research, and Western Washington University's Fund for the Enhancement of Graduate Research. I would like to thank my fellow graduate students, thesis committee, and adviser for academic support and feedback in all stages of my project. Much thanks for the support of my friends, family, and coworkers.

Capri!

TABLE OF CONTENTS

Abstract	iv
Acknowledgements	vi
Table of Figures	viii
Table of Tables	ix
Introduction.....	1
Materials and Methods.....	6
Results.....	17
Discussion.....	35
Literature Cited	49

TABLE OF FIGURES

Figure 1: Arrangement of diets (treatments) in 6-well plates	8
Figure 2: Radula tooth types and measurements	15
Figure 3: Placement of radula landmarks	16
Figure 4: Effect of diatom diet on the survival of postlarval <i>H. kamtschatkana</i>	18
Figure 5: Effect of diatom diet on the growth of postlarval <i>H. kamtschatkana</i>	23
Figure 6: Radula morphology examples: scanning electron microscope (SEM) images	28
Figure 7: Radula measurements as functions of shell length in 61-day old postlarvae	30
Figure 8: Average radula landmark positions by age	32
Figure 9: Vector description of the lack of difference between radula landmarks in postlarvae fed two different diets	34

TABLE OF TABLES

Table 1: Growth model selection table	20
Table 2: Number of surviving abalone per well does not affect growth	21
Table 3: Success of postlarval <i>H. kamtschatkana</i> on different diatom diets, and diatom characteristics.....	25
Table 4: Radula measurements by age.....	27
Table 5: Regression equations and r^2 values for radula morphology as a function of shell length	29
Table 6: Summary of literature on diatoms as food for abalone postlarvae	37

INTRODUCTION

Haliotis kamtschatkana kamtschatkana Jonas (pinto or northern abalone, hereafter referred to as *H. kamtschatkana*) ranges along the west coast of North America from Point Conception, California to southern Alaska, and is the only abalone found north of Oregon (Paul and Paul 1981; Geiger 2000). Combinations of poaching, commercial fishing, and recreational overharvesting severely diminished all populations of the species (Campbell 1999; Wallace 1999; Rothaus et al. 2008; Neuman et al. 2018; Carson and Ulrich 2019). Since abalone aggregate and broadcast spawn, adult density must be sufficiently high for gamete densities that allow fertilization (Allee 1949; Rothaus et al. 2008). Despite over two decades of harvest closure, persistently low adult *H. kamtschatkana* densities in the Salish Sea have caused reproduction and recruitment failure, leading to overall population decline that is unlikely to be reversed without human intervention (Rothaus et al. 2008; Bouma et al. 2012; Carson and Ulrich 2019).

Haliotis kamtschatkana restoration efforts in Washington State currently depend on hatchery production of juveniles by Puget Sound Restoration Fund. Restoration workers condition and spawn broodstock, rear larvae, induce settlement, and grow postlarvae first on diatom films, then on macroalgae (Vadopalas and Watson 2014). Juveniles are released at subtidal sites in the San Juan Archipelago when they are 8 to 45 mm in shell length (Vadopalas and Watson 2014). Hatchery workers have reported that low survival and slow growth make the early postlarval stage a bottleneck in efficient hatchery production of *H. kamtschatkana* (2015 conversation with Joshua V Bouma, unreferenced). Most mortality in abalones occurs in the first month post-settlement (Ebert and Houk 1984) and, anecdotally, *H. kamtschatkana* is more negatively affected by handling and experiences lower survival generally compared to other abalone species

(2015 conversation with Joshua V Bouma, unreferenced). Research specific to *H. kamtschatkana* postlarval cultivation is limited. Caldwell (1981) reported that “the overall survival [of *H. kamtschatkana*] would not lend itself well to a commercial operation”, and qualitatively described *H. kamtschatkana* postlarval growth rate as two-thirds that of *H. rufescens* Swainson, with high variability among individuals.

Abalone postlarvae begin to graze on benthic diatoms shortly after settlement, and the success of postlarvae depends on various diatom characteristics (Roberts, Kawamura, and Nicholson 1999; Gordon et al. 2006; Xing et al. 2007; Correa-Reyes et al. 2009). Depending on their size, postlarvae may graze on diatoms’ polysaccharide-based extracellular mucus (Hoagland et al. 1993; Kawamura and Takami 1995), consume whole diatoms (Kawamura, Roberts, and Nicholson 1998; Roberts, Kawamura, and Nicholson 1999), or rupture diatoms and consume the contents (Kawamura and Takami 1995). As postlarvae age and grow, they can consume different diatoms because the radula has larger teeth with steeper clearance angles that can break open diatoms that are tightly adhered or that have strong frustules (Kawamura, Roberts, and Nicholson 1998; Kawamura et al. 2001). The concurrent development of the digestive glands and stomach may also contribute to an increased ability to digest different diatoms (Takami et al. 1998).

Several factors influence which diatoms should be used in a hatchery to optimize abalone growth and survival. Diatom nutritional value, ingestibility, and growth rate depend on age of diatom film (Kawamura, Roberts, and Takami 1998), light intensity (Searcy-Bernal and Gorrostieta-Hurtado 2007), water flow (Searcy-Bernal and Gorrostieta-Hurtado 2007), inoculum density (Courtois de Vicose, Porta, et al. 2012), species of diatom (Roberts, Kawamura, and Nicholson 1999; Daume et al. 2000; Gordon et al. 2006; Xing et al. 2007; Correa-Reyes et al.

2009), and biochemical composition (Gordon et al. 2006; Correa-Reyes et al. 2009; Courtois de Vicoise, Porta, et al. 2012). Lacking research on the dietary value of different diatoms fed to *H. kamtschatkana* postlarvae, restoration hatcheries of this species have two choices: feed natural diatom film, which varies dramatically in quality and composition, or feed cultivated diatoms based on research of other abalone species. The latter can be problematic because *H. kamtschatkana* is found in cooler waters than other abalone species (Paul and Paul 1981; Paul and Paul 1998; Bouma 2007), and temperature affects diatom growth and survival, as well as the natural diatom communities in which *H. kamtschatkana* evolved. Thus, a diatom that works well for cultivation of one abalone species may not be appropriate for *H. kamtschatkana*.

The radula is important to postlarval success, because it affects the efficiency of food consumption, and improves digestion if it ruptures diatoms during consumption (Kawamura et al. 1995). Abalone have rhipidoglossan radulae, which are flexible and have many outer marginal teeth that sweep the substratum like a broom, in addition to a small number of more sturdy teeth in the center of each row (the rachidian tooth and lateral teeth; Steneck and Watling 1982; Ponder and Lindberg 1997). When postlarvae graze, the radula interacts directly with the diatoms, so radula morphology might be an important factor in diatom ingestibility (Roberts et al. 2001). Radula development has been characterized in several species of abalone (Roberts, Kawamura, and Takami 1999; Kawamura et al. 2001; Kawamura et al. 2001; Onitsuka et al. 2004; Johnston et al. 2005), but has not been examined in *H. kamtschatkana*. Differences in tooth number and rate of development are apparent between abalone species, but the basic process of development is the same: rows of teeth are formed at the posterior end of the radula, and worn teeth are shed from the anterior end (Moss 1999; Roberts, Kawamura, and Takami 1999; Kawamura et al. 2001; Takami et al. 2003; Onitsuka et al. 2004; Takami et al. 2006). As

abalone age, the number of rows per radula increases, the number of marginal and lateral teeth per row increases, and the shapes of the teeth change. An understanding of *H. kamtschatkana* radula development might indicate changes in the radula commensurate with changes in diatom diet as abalone grow, e.g. changes in tooth angle correlate with an ability to consume larger and more adhesive diatoms (Takami and Kawamura 2003) and teeth that are more numerous, more blunt, and larger correlate with the transition to macroalgae (Takami et al. 2003; Onitsuka et al. 2004). In addition, some species of Littorinidae have demonstrated radula plasticity in response to diatom diet type, but this has not been investigated in Haliotidae.

Abalone produce lecithotrophic larvae, which survive on energy reserves from egg yolk during development. These energy reserves are also available to postlarvae, demonstrated by the fact that starved postlarvae can survive for 12 days (*H. discus hannai*; Takami et al. 2000) or for 29 days (*H. iris*; Fukazawa et al. 2005), albeit with slowed growth (Takami et al. 2000; Roberts et al. 2001). Thus, even though diatom species identity (Kawamura, Roberts, and Takami 1998; Courtois de Vicose, Viera, et al. 2012) and density (Gorrostieta-Hurtado and Searcy-Bernal 2004) affect postlarval growth and survival, these changes might be undetectable until two weeks post-settlement if growing postlarvae are also relying on yolk reserves. There are no yolk reserve studies on *H. kamtschatkana* postlarvae, but we assume that this species is similar to others in that regard.

In the present study we fed *H. kamtschatkana* postlarvae one of 6 diatom species and measured survival, growth, and radula morphology at intervals over 61 days post-settlement. The species of diatoms were *Achnanthes brevipes*, *Amphora salina*, *Amphiprora paludosa*, *Cylindrotheca closterium*, *Navicula incerta*, and *Nitzschia laevis*, in addition to a starvation control. The survival and growth component of this project aimed to provide useful information

for *H. kamtschatkana* hatcheries when choosing species of diatoms as feeds. The radula component of this project aimed primarily to describe the morphology of the postlarval radula in *H. kamtschatkana* in the context of diet suitability and nutrient access, and secondarily to observe whether diatom diet can induce radula plasticity in the postlarval stage.

MATERIALS AND METHODS

We reared *H. kamtschatkana* postlarvae for 61 days post-settlement and fed them *ad libitum* on one of 6 species of benthic diatoms or in a starvation control. We counted survival at days 20, 26, 38, 49, and 61 post-settlement, measured growth at days 7, 20, 42, and 61 post-settlement, and collected postlarvae for radula analysis every other week. To investigate one basic component of nutrition, we measured the carbon-nitrogen ratio of the diatoms. We dissected radulae from animals, imaged them using scanning electron microscopy, and measured radula and tooth sizes and positions from the images. We investigated radula morphology over time and by size, and plastic response of radula morphology to diatom diet type.

Abalone for feeding trials

We obtained *H. kamtschatkana* larvae at 7 days post-fertilization from the Puget Sound Restoration Fund hatchery in Mukilteo, Washington. We transported them in a 4 L glass jar of 1 μm filtered seawater (FSW) within a cooler to Shannon Point Marine Center, Anacortes, Washington. Larvae were in transit for 3.5 hours and upon arrival at the marine center we allowed them to acclimate to the experimental incubator for 20 hours (12 °C, 12:12 light cycle, illuminated by four cool white fluorescent bulbs). We used UV-sterilized 0.2 μm FSW for all rinsing and rearing after larvae arrived at the laboratory.

To prepare the abalone for the feeding trial, we drained larvae onto a 60 μm Nitex mesh screen and rinsed them with FSW into a 1 L plastic beaker of 500 ml FSW. We then pipetted exactly 10 larvae into each well of 21 6-well culture plates and added FSW to a final volume of 14 ml. We arranged all diatom diets (treatments) such that every 7 wells contained a full set of all diets, randomly arranged (Figure 1). Thus, our experimental unit was a well and we had 18

wells per diatom diet. To induce the larvae to settle and undergo metamorphosis, we added gamma aminobutyric acid (GABA) to each well to a final concentration of 8 μ M (Paul Pratt, personal communication, unpublished manuscript). After 104 min (SD, 19 min), we conducted two 85% water changes in succession to remove the GABA, for a final volume of 10 ml of seawater in each well.




Figure 1: Arrangement of diets (treatments) in 6-well plates. Plate ID letter is at the top of each plate. Each row of plates in this figure was on a different shelf of the rearing incubator. Species are labeled for the first diet set, indicating the color coding of this figure. Abbreviations: A brev = *Achnanthes brevipes*, A pal = *Amphiprora paludosa*, A sal = *Amphora salina*, C clos = *Cylindrotheca closterium*, N lae = *Nitzschia laevis*, N inc = *Navicula incerta*, Starv = starvation control.

Diatoms for feeding trials

We purchased 6 species of benthic diatoms: *Achnanthes brevipes* Agardh (CCMP 100), *Amphiprora paludosa* Van Heurck (CCMP 125; synonym: *Entomoneis paludosa*), *Amphora salina* W. Smith (CCMP 1119; synonyms: *Amphora coffeaeformis*, *Halamphora coyfefe*, *Halamphora coffeaeformis*), *Cylindrotheca closterium* (Ehr.) Reimann et Lewin (CCMP 340; synonym: *Nitzschia closterium*), *Navicula incerta* Grunow (CCMP 542), and *Nitzschia laevis* Hustedt (CCMP 559), from the National Center for Marine Algae and Microbiota (Bigelow Laboratory, East Boothbay, Maine). We chose these species based on size, shape, preferred temperature range, and information from other feeding studies on abalone postlarvae.

Prior to inoculation into wells containing abalone postlarvae, we cultured the diatoms at 17 °C on a 16:8 hour light:dark cycle in axenic conditions in 1 L flasks of approximately 600 ml Proline F/2 media with added sodium metasilicate (Aquatic Eco-Systems Inc., Apopka, FL). Sixteen hours after the abalone settled, we added diatoms at 1 cell mm⁻² of wetted well surface area (2,200 cells per well) to the wells, held at 12 °C on a 12:12 hour light:dark cycle. Thereafter, we visually observed that diatoms grew at a rate greater than or equal to postlarval grazing rates and needed no further supplementation. We conducted 80% FSW changes 3 times per week in a 12 °C cold room with 5 ml glass pipets. When diatoms appeared to be growing more than one cell deep, at day 45, we thinned them by brushing well bottoms with fringed waterproof paper.

To determine C:N ratios, we allowed diatoms to grow in wells in the same conditions for one week after postlarvae were removed on day 61. During this week, we conducted water changes as previously described. We then dislodged diatoms and stirred them into suspension in the wells using strips of silicone cut from a 3/8" hose. We pipetted the diatom suspension into a syringe

fitted with a 0.7 μm glass fiber filter and placed the filters into tin capsules held in a 24-well culture plate. We dried the filters at 50 °C for 36 hours, then folded the tin capsules closed. Samples were analyzed for ^{13}C and ^{15}N isotopes at the University of California Davis Stable Isotope Laboratory (Davis, California), using an Elementar Micro Cube elemental analyzer interfaced to an Isoprime VisION IRMS (Elementar Analysensysteme GmbH, Hanau, Germany).

We determined mean C:N ratio from each diatom diet and related it qualitatively to survival and growth of postlarvae fed the different diatom diets.

Survival

We counted the number of abalone alive in each well at days 20, 26, 38, 49, and 61 post-settlement. Prior to each count, we used a 1 ml micropipetor to flush water across each postlarva to dislodge the dead and leave the live postlarvae in place.

Because most mortality occurred before our first measurement of survival on day 20, we were unable to fit a Survival Analysis curve. Instead we analyzed percent survival at day 20 and day 61. For both periods, we tested for effect of diatom diet using a one-way ANOVA, followed by pairwise contrasts (Tukey HSD) to compare individual diets to each other. We omitted the starvation control and *A. brevipes* from the day 61 analysis because of extremely low survival (see Results). Since there were no *a priori* hypotheses about how diatom diets would compare to each other, we used pairwise contrasts rather than special contrasts. We chose the Tukey HSD test because it is widely used, we wanted a more sensitive test than a Bonferroni, and we had too many comparisons to use a Fischer's LSD test.

We checked for homogeneity of variance using Levene's test. Variance of survival data was homogenous ($p = 0.55$ for day 20; $p = 0.11$ for day 61). We checked for normality using the

Shapiro-Wilk test with a threshold of $p \geq 0.01$. Two treatments did not meet this normality threshold (*A. brevipes*, $p = 0.0097$ for day 20; *C. closterium* $p = 0.007$ for day 61). To reduce the risk of false positives due to non-normality, we chose an α level of 0.01 for our Tukey HSD test.

Growth

We measured growth of postlarval abalone at days 7, 19, 20, 42, and 61 post-settlement. Day 7 measurements were only of postlarvae residing on the well bottoms, because animals on the well sides were too delicate at this age to dislodge for measurement. Beyond day 7, postlarvae typically resided on the walls of wells and were rarely observed on the bottoms, but were robust enough to be moved to the bottom of the well for measurement. We dislodged all postlarvae in each well using thin grass stems, then manipulated them onto their foot on the bottom center of the well. We dipped the stems in boiling water to prevent diatom contamination between wells. We photographed the postlarvae using a stereoscope equipped with a camera (optical lens magnification 8 \times ; Leica Microsystems, Buffalo Grove, IL). We then measured shell length using ImageJ image analysis software (Rasband 1997).

To compare growth among diatom diets, we used a linear mixed model fit by maximum likelihood and with effective degrees of freedom calculated by Satterthwaite's method (Bates et al. 2015; Kuznetsova et al. 2017). Linear mixed models are appropriate for data that include repeated measurements and have missing data. We calculated the mean shell length of all postlarvae in each well on each measurement day, then natural-log transformed the shell length data so that it fit a linear relationship with age. We first fit the full model, containing the fixed factors diatom diet, age, and diatom diet-age interaction, and the random factors diet set, culture plate, and well (experimental unit). Well was required as a random factor because each well was

measured repeatedly throughout the experiment. We then compared the full model to reduced models using the Akaike information criteria and an ANOVA.

After developing the growth models for diets, we tested whether size of postlarvae was influenced by the number of surviving postlarvae in a well, e.g. by competition and crowding. We used data from the four diets with highest survival, because the data were more robust. For each of these diets, we fitted a linear mixed model containing the fixed factors age and count, and the random factor well (experimental unit). Age was transformed as $e^{0.02*\text{day}}$, based on our knowledge from the diet-growth model results, and was required in the model because shell length is highly dependent upon age. Well was required as a random factor because each well was measured repeatedly throughout the experiment.

Radula morphometry

During the feeding trial, we removed one diet set of postlarvae at days 5, 20, and 33 post-settlement, and preserved postlarvae in deionized water in a -20 °C freezer for radula analysis. At day 61, we preserved all surviving postlarvae from the feeding trial using the same methods. In addition, we obtained a few juvenile (9 month old) animals from Puget Sound Restoration Fund for radula dissection and comparison. We extracted each radula by thawing the abalone, then adding 6% sodium hypochlorite to dissolve soft tissues. We used a pipet to flush the radula free of remaining soft tissues, then rinsed it thoroughly with deionized water. Once each radula was rinsed, we placed it on a thin plastic sheet in a droplet of water. Once the radula was dry, we transferred it to a scanning electron microscope aluminum stub by touching it gently with double-stick carbon tape.

We coated the samples with gold-palladium sputter (60 seconds, 26 kV; Polaron Range; Quorum Technologies, East Sussex, England) then used a scanning electron microscope (Vega TS 5136MM, TESCAN, Kohoutovice, Czech Republic) to obtain an image of each whole radula at 1,000× and an image of 5 to 10 rows in the middle of each radula at 2,000× to 6,000×.

We conducted two types of morphometric analysis: traditional and geometric. For traditional morphometric analysis, we measured radula length, radula width, gap length (distance between rows measured from rachidian base to rachidian base), number of rows, buccal cartilage position (row number at which the posterior ends of buccal cartilages terminate), rachidian height (base to fold), rachidian cusp length (fold to tip), rachidian base width, number of lateral teeth per row, and number of marginal teeth per row (Figure 2). We plotted these measurements against shell length, grouped by diatom diet, following procedures in Avaca et al. (2010). We fit simple linear regressions of traditional morphometric measurements predicted by shell length. We described tooth development with age (rather than with size) qualitatively, because of the limited numbers of postlarvae sampled earlier than 61 days post-settlement.

We observed no difference in radula morphology between diets or wells, and had a limited number of successful dissections, so we treated radulae as individual units, rather than averaging characteristics within wells. For all radula measurements except for radula length, width, and buccal cartilage position, we measured multiple rows or rachidian teeth and averaged them into a single datum per radula.

Our geometric morphometric analysis was based on 2-dimensional landmarks. We described overall shape of rachidian teeth using 7 landmarks, and used 13 landmarks to describe the relative positions of the middle 5 teeth within a row: the rachidian tooth and the first 2 lateral

teeth (L1 and L2) on either side of it (Figure 3). We aligned and scaled sets of landmarks using Procrustes General Analysis, which centers, scales, and then rotates shapes until the sum of squared distances among them is minimized. We estimated positions of missing landmarks using multivariate regression estimates (Adams et al. 2019), then averaged the coordinates of landmarks from different rows within each radula. Bilateral symmetry was not assumed for abalone radulae (Hickman 1981; Geiger 1999). We then compared radula landmarks between diatom diets using Procrustes ANOVA (Adams et al. 2019).




Figure 2: Radula tooth types and measurements. Arrow at left indicates base of row where buccal cartilage ends, as measured in this study.





Figure 3: Placement of radula landmarks used to analyze radula morphology. Above: rachidian: 7 landmarks (n = 394 rachidian teeth from 78 radulae; 76 missing landmarks from 49 radulae were estimated). Below: the 5 center teeth: L2-L1-rachidian-L1-L2: 13 landmarks (n = 215 rows from 86 radulae; 64 missing landmarks from 45 radulae were estimated).

RESULTS

Survival

Amphora salina yielded the highest survival (Tukey HSD, $p < 0.03$), in a statistically homogenous subgroup with *N. incerta*, *A. paludosa*, and *C. closterium*. *Navicula incerta*, *A. paludosa*, *C. closterium*, *N. laevis*, *A. brevipes*, and starvation formed a second homogenous subgroup (Tukey HSD $p > 0.05$, Figure 4). *Nitzschia laevis* yielded mediocre survival at day 20, and poor survival at day 61. *Achnanthes brevipes* yielded survival lower than starvation at day 20, and very poor survival at day 61 (Figure 4).

Diatom diet had a significant effect on postlarval survival at both day 20 (ANOVA, $F_{6,95} = 8.60$, $p < 0.001$) and day 61 (ANOVA, $F_{6,66} = 9.70$, $p < 0.001$). At our first measurement of survival, on day 20, all diets yielded less than 40% survival, except *A. salina* with 60%. We did not measure settlement or metamorphosis success separately from survival, so both larvae that failed to settle and postlarvae that died after settlement reduced the percent survival. Therefore, survival at day 20 may be due to the diatoms' suitability as a feed very early in life, suitability for settlement and metamorphosis prior to feeding, or a combination.

For postlarvae fed *N. incerta*, *A. salina*, *A. paludosa*, or *C. closterium*, survival increased substantially after day 20. Of postlarvae fed these diets and alive on day 20, 78% were still alive on day 61 (Figure 4). Differences in survival between these diatom diets from days 0 to 61 were driven by early survival to day 20, indicating that either it is a refuge age of higher survival, or that these diets are poorly suitable for settlement and metamorphosis, thus causing early death. Our last observation of a living starved postlarva was on day 26.




Figure 4: Effect of diatom diet on the survival of postlarval *H. kamtschatkana*. Error bars represent 90% confidence interval. Letters below each bar represent statistically homogeneous sub-groups (ANOVA $p > 0.05$): letters a and b for day 20; letters y and z for day 61.

Growth model selection

The most parsimonious linear mixed model to predict growth of postlarvae included diatom diet, postlarval age, and diet-age interaction as fixed predictive factors, and well (experimental unit) as a random factor with both a random slope and a random intercept (Table 1). Random factors of culture plate and diet set added little to the models and were left out of the final model (Table 1).

There was a significant difference in modeled initial shell length (intercept) of postlarvae fed *A. salina* compared to starting size of postlarvae fed other diets, even though the actual initial shell length was not different between diets. We viewed this as an unavoidable and insignificant artifact of model fitting.

Survival rates of postlarvae fed *A. brevipes* or the starvation control were too low for statistical analysis of growth, so they were left out of the analysis. Number of surviving postlarvae per well did not affect subsequent growth, indicating that there were no density effects, and space and food were not limiting resources (Table 2).

Table 1: Growth model selection table. All models include a random slope across all ages for each random factor. Diet = diatom treatment. Age = days post-settlement. Well and plate = growing containers. † no significant difference between models, ANOVA $p > 0.05$

Model #	Predictive factors included in model:	Factors for which a random slope is included across all ages:	df	AIC
† 1)	diet, age, diet-age interaction	well, plate	21	-478
† 2)	diet, age, diet-age interaction	well	18	-477
† 3)	diet, age, diet-age interaction	well, diet set	21	-473
† 4)	diet, age, diet-age interaction	well, diet set, plate	24	-472
5)	diet, age	well, plate	15	-468
6)	diet, age	well	12	-467
7)	diet, age	well, diet set	15	-464
8)	diet, age	well, diet set, plate	18	-462
9)	age	well	6	-442
10)	diet, age, diet-age interaction	plate	18	-418
11)	diet, age, diet-age interaction	diet set	18	-413
12)	diet, age, diet-age interaction	diet set, plate	21	-412
13)	diet, age	plate	12	-389
14)	diet, age	diet set	12	-386
15)	diet, age	diet set, plate	15	-383

Table 2: Number of surviving abalone per well does not affect growth. Data are values for shell length within each diet, described by the equation $SL = \beta_0 + \beta_{age}A + \beta_{number}N + e_{well} + \varepsilon_i$, where SL is shell length, A is postlarval age, N is number of postlarvae per well, e is the random effect of well, and ε_i is error. df = Satterthwaite's effective degrees of freedom.

Diet	Intercept	β_{age}	β_{number}	Statistics for β_{number}		
				df	t-value	p-value
<i>A. paludosa</i>	25	275	0.15	53	0.045	0.97
<i>A. salina</i>	228	218	-10	57	-1.7	0.09
<i>C. closterium</i>	-56	347	0.99	43	0.16	0.87
<i>N. incerta</i>	103	237	-2.4	43	-0.87	0.39

Growth of postlarvae

Between most diets, there was no significant difference in growth of postlarvae. The exception to this was *C. closterium*, which yielded significantly faster growth than *A. salina* or *N. incerta* (LMM $p = 0.001$ and 0.018 , Satterthwaite effective $df = 59$ and 64 , respectively), and nearly significantly faster than those fed *A. paludosa* or *N. laevis* (LMM $p = 0.075$ and 0.088 , Satterthwaite effective $df = 61$ and 70 , respectively; Figure 5).

The growth of *H. kamtschatkana* postlarvae when fed *C. closterium* is described by the equation

$$\text{shell length } (\mu\text{m}) = 293 * e^{0.021t}$$

where $t =$ days post-settlement.

There was no significant difference in growth rate between postlarvae fed *A. paludosa*, *A. salina*, *N. incerta*, or *N. laevis* (LMM $p > 0.10$, Satterthwaite effective df range = 63 to 75). The growth of postlarvae fed these diatom diets was:

$$\text{shell length } (\mu\text{m}) = 302 * e^{0.018t}$$

By day 61, mean shell length among all postlarvae regardless of diet was $944 \mu\text{m}$ (SD, $186 \mu\text{m}$; range of individual postlarvae: 558 to $1,390 \mu\text{m}$), and none of the postlarvae developed a respiratory pore within the course of our experiment.




Figure 5: Effect of diatom diet on the growth of postlarval *H. kamtschatkana*. Error bars represent one standard error of the mean. Curves are of the linear mixed model of shell length in response to postlarval age, diatom diet, and age-diet interaction, with well (experimental unit) as a random factor.

Diatom characteristics and carbon:nitrogen ratio

We measured C:N ratio for all diatoms, measured length and width of 4 of the 6 diatoms, and noted growth patterns and shape. C:N values had low variance within diatom diets (Table 3), and there was no clear relationship between C:N ratio and either survival or growth of postlarvae. Likewise, there were no apparent relationships between diatom size and survival or growth of postlarvae

We observed that *A. paludosa* adhered very loosely to container surfaces, almost like a planktonic diatom. *Nitzschia laevis* tended to clump, while *A. salina*, *C. closterium*, and *N. incerta* formed uniform films. *Achnanthes brevipes* and *N. laevis* appeared to grow more slowly than the other diatoms tested, but all diatoms used in this study grew well at 12 °C. This temperature was within the reported range of all diatom diets except for *C. closterium*, which had a reported temperature range of 18 to 26 °C (NCMA 2019a).

Table 3: Success of postlarval *H. kamtschatkana* on different diatom diets, and diatom characteristics. Diatom sizes are as measured in the present study and as reported by the Bigelow Laboratory (NCMA 2019b). One standard deviation is given after \pm ; nd = no data. Sample size for diatom length and width is >20 per diatom; n refers to number of wells that were analyzed for C:N ratio.

Diatom species	Length (μm)	Bigelow length (μm)	Width (μm)	Bigelow width (μm)	Survival	Growth	C:N	n
<i>A. salina</i>	nd	24 - 30	nd	6 - 8	Excellent	Fair	30 ± 2	9
<i>N. incerta</i>	18	12 - 16	7.9	4 - 6	Good	Fair	24 ± 3	8
<i>C. closterium</i>	13	18 - 22	4.2	2 - 3	Fair	Excellent	10 ± 2	10
<i>A. paludosa</i>	13	9 - 21	7.9	6 - 9	Fair	Good	10 ± 4	8
<i>N. laevis</i>	15	16 - 20	6.5	4 - 6	Poor	Good	21 ± 3	6
<i>A. brevipes</i>	nd	6 - 20	nd	6 - 13	Poor	Poor/NA	22	1

Radula development with age and size: traditional morphometrics

As postlarvae aged, we observed an increase in rachidian tooth size, radula length and width, number of teeth per row, and number of rows per radula (Table 4). For all ages and sizes of postlarvae in our study, qualitative tooth shape differences were small: tooth angles and shapes were very similar among all postlarval radulae, and the prominent difference was in the size and number of teeth rather than the shape (Figure 6). From day 5 to day 61, the radula gained an average of 10 marginal teeth and 2.7 lateral teeth. Radula length approximately tripled in this time from 68 to 220 μm , and radula width approximately doubled from 17 to 41 μm (Table 4).

In 61-day old postlarvae, shell length was a linear predictor of radula size, rachidian tooth size, gap length, and number of both marginal and lateral teeth per row (Figure 7, Table 5).

The buccal cartilage position did not change with age or shell length, indicating that postlarvae graze using 10 rows (SD, 1.5 rows) of teeth regardless of postlarval age or size. Number of rows per radula increased slightly with age but not with day 61 shell length.

Table 4: Radula measurements by age. Data include all diatom diets. Number of lateral teeth are total per row of the radula; number of marginal teeth are per one side of row. Number of rows are per radula. Buccal cartilage position refers to the number of rows anterior to and adjoining the buccal cartilages; in other words, the row number where the cartilages end. Rach = rachidian tooth. One standard deviation is given after \pm ; nd = no data. Sample sizes are of individual postlarvae, shown in parentheses. *This shell length is of the study population, because we did not measure individuals prior to dissection before day 61.

Age (d)	Shell length (μm)	No. lateral teeth	No. marginal teeth	No. rows	Buccal cartilage position	Radula length (μm)	Radula width (μm)	Gap length (μm)	Rach. height (μm)	Rach. tip (μm)
5	nd	2	2	17	nd	68	17	4.5	3.8	1.8
		(2)	(2)	(1)		(1)	(4)	(4)	(2)	(3)
20	440 $\pm 52^*$	2	4	21	10	104	22	4.8	3.9	2.2
		(11)	(11)	(9)	(10)	(9)	(11)	(11)	(9)	(9)
33	nd	4	7	23	11	139	28	5.9	5.0	2.7
		(14)	(12)	(13)	(14)	(14)	(16)	(15)	(14)	(14)
61	903 ± 179	5	12	23	10	221	41	9.0	6.7	3.2
		(100)	(91)	(83)	(65)	(80)	(94)	(111)	(98)	(98)




Figure 6: Radula morphology examples: scanning electron microscope (SEM) images of radulae at days 5, 19, 33, and 61 post-settlement, and 9-month old juvenile. Top-right image shows tooth-down view of the radula. Note buccal cartilages in top left images (arrows). Scale bars represent 10 μm , except 100 μm for 9-month old juvenile radula.

Table 5: Regression equations and r^2 values for radula morphology as a function of shell length. The independent variable (x) is shell length (mm). Sample sizes are of individual postlarvae.

Variable	Slope	Intercept	r^2	n
Gap length (μm)	9.0	1.0	0.93	106
Radula width (μm)	42	3	0.93	89
Rachidian height (μm)	6.2	1.1	0.91	93
Radula length (μm)	234	9	0.90	77
Marginal teeth per row	8.4	4.5	0.88	86
Rachidian tip (μm)	2.1	1.3	0.81	93
Lateral teeth per row	3.6	1.6	0.76	95
Rows per radula	-0.6	23	0.19	79
Buccal cartilage position	2.0	8.3	0.01	62
Rachidian ratio	0.4	1.8	0.01	93







Figure 7: Radula measurements as functions of shell length in 61-day old postlarvae. Data include all diatom diets. Each data point represents one postlarva. Linear regression lines were calculated using the least squares method; see Table 5 for r^2 values and equations of the lines.

Radula development with age and size: geometric morphometrics

In addition to traditional morphometric measurements, we analyzed landmark positions of rachidian teeth and of the middle 5 teeth of each radula (Figure 3). Our rachidian tooth landmarks marked the relative tip position, which could indicate tooth contact angle with the grazing surface, and marked three different widths of the hourglass-shaped rachidian teeth, which could reveal changes in overall structure. There were no apparent changes of position of any of these landmarks (Figure 8). Although the number of teeth in each row increased with size and age, the positions of landmarks on the middle 5 teeth did not change relative to each other with age (Figure 8).




Figure 8: Mean radula landmark positions by age. Colored markers represent landmark positions (Figure 3) at different ages. Landmarks were centered, scaled, and aligned by Procrustes General Analysis. The gray landmark dots and tooth outlines are examples to assist the reader in understanding the landmarks, and not representative of any one result.

Lack of radula plasticity in response to diet

The diatom diets we tested had no effect on any traditional morphometric or landmark measurement of radula morphology. Orientations of landmarks of the rachidian tooth, and landmarks of the middle 5 teeth were not significantly different between diatom diets (Procrustes ANOVA for rachidian $F_{4,97} = 0.68$; $p = 0.8$; for row $F_{4,78} = 1.1$; $p = 0.4$).

Postlarvae fed *C. closterium* and *N. laevis* showed more difference in row-center landmarks than any other pair of diets (Figure 9). Even so, differences between these two diets are so small that many position-change arrows are obscured by the reference points: where two marks exit the reference point, these two marks are the back sides (barbs) of the arrow.




Figure 9: Vector description of the lack of difference between radula landmarks in postlarvae fed two different diets: *C. closterium* (reference points) vs. *N. laevis* (tips of arrows). This is the greatest difference in tooth morphology that we found, indicating a lack of plastic response to diet. Note that some reference points obscure the arrow tips and only the arrow “barbs” are visible. The gray landmark dots and tooth outlines are examples to assist the reader in understanding the landmarks, and not representative of any one result.

DISCUSSION

Effect of diatom diet

Amphora salina was the best diet for survival of postlarval *H. kamtschatkana*. It appears to be a good diet for other abalone species as well, although it appears only twice in abalone feeding literature to date (Table 6): 77% of *H. asinina* Linnaeus fed *A. salina* survived to day 20 post-settlement (Ding et al. 2017), and *H. diversicolor supertexta* postlarvae preferentially consumed *A. salina* from natural mixed diatom films (Zhang et al. 2010). *Amphora salina* grows singly (Wang et al. 2014) or in pairs. Its organic layer is smooth and tight on the silica frustule, and there is an additional glucose polymer present between cells after division (Tesson and Hildebrand 2013), which may contribute to extracellular nutritional availability to small postlarvae. Extracellular polymeric substances may affect food preferences, digestibility, and particle sizes consumed (Joyce and Utting 2015). This could help explain the success of *A. salina* as a diet in our study, in spite of it being the largest of the diatoms that we tested.

Cylindrotheca closterium was the best diet for growth of postlarval *H. kamtschatkana* in our study. *Cylindrotheca closterium* is very well represented in postlarval feeding literature as a successful feed for a variety of abalone species. Its frustule is thin (Reimann and Lewin 1964); this may increase nutritional value because the cell contents can be digested if the frustule breaks during feeding. Postlarvae of *H. rufescens*, and likely other abalone species as well, will pass whole, live diatoms through the digestive system if the diatom frustule is strong enough to remain intact during feeding (Argumedo-Hernández et al. 2010).

Many abalone feeding papers claim *C. closterium* has weak adhesion, but the basis for this traces back to only one paper, which did not directly measure adhesion (Kawamura and Hirano 1992). In our observation, *C. closterium* did not have particularly weak adhesion and instead

formed a substantial film that did not dislodge when culture flasks were swirled. Adhesive properties of benthic diatoms depend not only on species, but also on strain, growing conditions (de Brouwer and Stal 2002; Ravizza and Hallegraeff 2015), and surface material (Rasmussen and Østgaard 2001; Holland et al. 2004), any of which might explain this discrepancy. Regardless of the adhesion level, *H. kamtschatkana* postlarvae appeared to have no difficulty consuming *C. closterium*. Near the end of the 9-week experiment we observed that the films of *C. closterium* were nearly grazed clean, and feces were numerous. This indicates a high level of consumption that would have approached food limitation had our experiment continued.

A combination of *A. salina* and *C. closterium* might be favorable to hatchery success, since the former best supported survival and the latter best supported growth, particularly after the first four weeks post-settlement. Diatoms fed in combination can successfully support postlarval settlement, growth, and survival in *H. rufescens* (Araya et al. 2010), *H. fulgens* Philippi (Viana et al. 2007), and *H. discus hannai* Ino (Gordon et al. 2006), but for simplicity most diatom feeding experiments test only single-species diets. When feeding two or more diatoms simultaneously, it is important that both grow well and neither out-competes the other. *Cylindrotheca closterium* has some allelopathic effects against dinoflagellates but these effects have not been shown against diatoms (Xu et al. 2019); for example, it grows viably with *Navicula* sp. (Najmudeen 2017). However, *C. closterium* grows especially quickly during initial surface colonization (Najmudeen 2017), so to co-culture our highest performing diets, *A. salina* must either grow equally quickly during initial surface colonization alongside *C. closterium*, or continue to grow robustly despite this initial competition. Another option is to inoculate *A. salina* in advance so that it can begin colonization in the absence of competition, then add *C. closterium* a few days later.

Table 6: Summary of literature on diatoms as food for abalone postlarvae of shell lengths less than 800 μm . Where growth and/or survival are given as a range, this is often because the experiment was testing other culture parameters (e.g. artificial lighting, water flow). Age refers to days post-settlement. Ages in parentheses are experimental durations, for references where initial age is not given. SL = shell length. Blank fields: information not given in reference.

Diatom	Abalone	Growth ($\mu\text{m d}^{-1}$)	Survival (%)	Initial SL (μm)	Initial Age (d)	Final Age (d)	Temp. ($^{\circ}\text{C}$)	Reference
<i>Achnanthes brevipes</i>	<i>H. discus hannai</i>	57		1,300	28	41	20	Kawamura et al. 1995
<i>Achnanthes longipes</i>	<i>H. discus hannai</i>	48		1,300	28	41	20	Kawamura et al. 1995
<i>Achnanthes longipes</i>	<i>H. discus hannai</i>	9	70	674		(7)	20	Takami et al. 2003
<i>Amphora angusta</i>	<i>H. discus hannai</i>	30		1,300	28	40	20	Kawamura et al. 1995
<i>Amphora luciae</i>	<i>H. discus hannai</i>	24	43	280	78	109	22	Gordon et al. 2006
<i>Amphora proteus</i>	<i>H. discus hannai</i>	12	35		19	38		Xing et al. 2007
<i>Amphora salina</i>	<i>H. discus hannai</i>	19	26		9	28		Xing et al. 2007
<i>Cocconeis scutellum</i>	<i>H. discus hannai</i>	8	100	350	11	25	20	Kawamura and Takami 1995
<i>Cocconeis scutellum</i>	<i>H. discus hannai</i>	25		350	0	10	20	Kawamura and Takami 1995
<i>Cocconeis scutellum</i>	<i>H. discus hannai</i>	41		1,300	28	40	20	Kawamura et al. 1995
<i>Cocconeis scutellum</i>	<i>H. discus hannai</i>		0	~350	0	28	20	Takami et al. 1997
<i>Cylindrotheca closterium</i>	<i>H. discus hannai</i>	22	63	350	11	25	20	Kawamura and Takami 1995
<i>Cylindrotheca closterium</i>	<i>H. discus hannai</i>	36		350	0	10	20	Kawamura and Takami 1995
<i>Cylindrotheca closterium</i>	<i>H. discus hannai</i>	50		1,300	28	39	20	Kawamura et al. 1995
<i>Cylindrotheca closterium</i>	<i>H. discus hannai</i>	21	90	447	25	32		Takami et al. 2003
<i>Cylindrotheca closterium</i>	<i>H. discus hannai</i>	33	88	657	27	34	20	Takami et al. 2003
<i>Cylindrotheca closterium</i>	<i>H. discus hannai</i>	44	78	854	21	29	20	Takami et al. 2003
<i>Hantzschia amphioys</i>	<i>H. discus hannai</i>	11	47		20	39		Xing et al. 2007
Mix: <i>Navicula lenzii</i> , <i>Amphora luciae</i>	<i>H. discus hannai</i>	36	49	280	2	33	22	Gordon et al. 2006
Mix: <i>Navicula lenzii</i> , <i>Nitzschia laevia</i>	<i>H. discus hannai</i>	23	42	280		(31)	22	Gordon et al. 2006

Diatom	Abalone	Growth ($\mu\text{m d}^{-1}$)	Survival (%)	Initial SL (μm)	Initial Age (d)	Final Age (d)	Temp. ($^{\circ}\text{C}$)	Reference
Mix: <i>Navicula lenzii</i> , <i>Nitzschia laevis</i> , <i>Amphora luciae</i>	<i>H. discus hannai</i>	33	40	280	36	67	22	Gordon et al. 2006
Mix: <i>Nitzschia laevis</i> , <i>Amphora luciae</i>	<i>H. discus hannai</i>	27	31	280	23	54	22	Gordon et al. 2006
<i>Navicula cf. lenzii</i>	<i>H. discus hannai</i>	21	40	280		(31)	22	Gordon et al. 2006
<i>Navicula corymbosa</i>	<i>H. discus hannai</i>	31	84		10	29		Xing et al. 2007
<i>Navicula parva</i>	<i>H. discus hannai</i>	12	74		32	51		Xing et al. 2007
<i>Navicula ramosissima</i>	<i>H. discus hannai</i>	16	67	350	11	25	20	Kawamura and Takami 1995
<i>Navicula ramosissima</i>	<i>H. discus hannai</i>	18		350	0	10	20	Kawamura and Takami 1995
<i>Navicula ramosissima</i>	<i>H. discus hannai</i>	23		1,300	28	41	20	Kawamura et al. 1995
<i>Navicula seminulum</i>	<i>H. discus hannai</i>	12	51			(19)		Xing et al. 2007
<i>Navicula</i> sp.	<i>H. discus hannai</i>	46	73	360		(40)	22	Pang et al. 2006
<i>Nitzschia laevis</i>	<i>H. discus hannai</i>		4	280		(31)	22	Gordon et al. 2006
<i>Nitzschia</i> sp.	<i>H. discus hannai</i>	14		1,300	28	40	20	Kawamura et al. 1995
<i>Nitzschia</i> sp.	<i>H. discus hannai</i>	35	95		46	65		Xing et al. 2007
<i>Pleurosigma</i> sp.	<i>H. discus hannai</i>	24		1,300	28	39	20	Kawamura et al. 1995
<i>Rhaphoneis surirella</i>	<i>H. discus hannai</i>	28	86		40	59		Xing et al. 2007
<i>Stauroneis constricta</i>	<i>H. discus hannai</i>	19	65	350	11	25	20	Kawamura and Takami 1995
<i>Stauroneis constricta</i>	<i>H. discus hannai</i>	26		350	0	10	20	Kawamura and Takami 1995
<i>Synedra investiens</i>	<i>H. discus hannai</i>	18		1,300	28	40	20	Kawamura et al. 1995
<i>Navicula incerta</i>	<i>H. fulgens</i>	15	>95	~250	0	15	18	Searcy-Bernal et al. 2001
<i>Navicula incerta</i>	<i>H. fulgens</i>	33	>95	474	15	60	18	Searcy-Bernal et al. 2001
<i>Cocconeis scutellum</i>	<i>H. iris</i>	45	70	650	24	92	18	Roberts, Kawamura, and Nicholson 1999
<i>Cocconeis</i> sp.	<i>H. iris</i>	33 - 42	~80	1,400		(14)	18	Uriarte et al. 2006
<i>Cylindrotheca closterium</i>	<i>H. iris</i>	40	75	650	33	101	18	Roberts, Kawamura, and Nicholson 1999
<i>Cylindrotheca closterium</i>	<i>H. iris</i>	42 - 48	~90	1,400		(14)	18	Uriarte et al. 2006

Diatom	Abalone	Growth ($\mu\text{m d}^{-1}$)	Survival (%)	Initial SL (μm)	Initial Age (d)	Final Age (d)	Temp. ($^{\circ}\text{C}$)	Reference
<i>Navicula britannica</i>	<i>H. iris</i>	25	48	650	28	96	18	Roberts, Kawamura, and Nicholson 1999
<i>Navicula ramosissima</i>	<i>H. iris</i>	25	100	650	35	68	18	Roberts, Kawamura, and Nicholson 1999
<i>Nitzschia ovalis</i>	<i>H. iris</i>	23 - 45	~90	1,400		(14)	18	Uriarte et al. 2006
<i>Pleurosigma</i> sp.	<i>H. iris</i>	1	0	650		(54)	18	Roberts, Kawamura, and Nicholson 1999
<i>Achnanthes brevipes</i>	<i>H. kamtschatkana</i>	4	1	332	7	61	12	present study
<i>Amphiprora paludosa</i>	<i>H. kamtschatkana</i>	11	26	339	7	61	12	present study
<i>Amphora salina</i>	<i>H. kamtschatkana</i>	10	47	349	7	61	12	present study
<i>Cylindrotheca closterium</i>	<i>H. kamtschatkana</i>	14	22	337	7	61	12	present study
<i>Navicula incerta</i>	<i>H. kamtschatkana</i>	10	37	334	7	61	12	present study
<i>Nitzschia laevis</i>	<i>H. kamtschatkana</i>	11	6	327	7	61	12	present study
<i>Cocconeis</i> sp.	<i>H. rubra</i>	27	71	400	12	87	17	Daume et al. 2000
<i>Cylindrotheca closterium</i>	<i>H. rubra</i>	33	17	400	44	121	17	Daume et al. 2000
<i>Navicula jeffreyi</i>	<i>H. rubra</i>	35	63	400	21	98	17	Daume et al. 2000
<i>Navicula</i> sp.	<i>H. rubra</i>	39	75	400		(77)	17	Daume et al. 2000
<i>Navicula</i> sp.	<i>H. rubra</i>	19	0 - 40	490		(42)	18	Day et al. 2004
<i>Navicula</i> sp.	<i>H. rubra</i>	12	20 - 85	663		(21)	18	Day et al. 2004
<i>Navicula</i> sp.	<i>H. rubra</i>	8	5 - 10	663	39	60	18	Day et al. 2004
<i>Amphiprora paludosa</i>	<i>H. rufescens</i>	26		~300	5	55	18	Correa-Reyes et al. 2009
<i>Navicula incerta</i>	<i>H. rufescens</i>	21		~300	5	55	18	Correa-Reyes et al. 2009
<i>Navicula incerta</i>	<i>H. rufescens</i>	48 - 62	27 - 68	490	9	72	18	Anguiano-Beltran and Searcy-Bernal 2013
<i>Navicula incerta</i>	<i>H. rufescens</i>	31 - 38	52 - 80	332	6	50	17	Searcy-Bernal & Gorrostieta-Hurtado 2007
<i>Navicula incerta</i>	<i>H. rufescens</i>	15 - 22	54 - 67	1,250		(30)	16	Uriarte et al. 2006
<i>Nitzschia laevis</i> strain "B"	<i>H. rufescens</i>	13		~300	5	55	18	Correa-Reyes et al. 2009
<i>Nitzschia laevis</i> strain "C"	<i>H. rufescens</i>	19		~300	5	55	18	Correa-Reyes et al. 2009
<i>Nitzschia laevis</i> strain "D"	<i>H. rufescens</i>	20		~300	5	55	18	Correa-Reyes et al. 2009
<i>Nitzschia</i> cf. <i>fonticola</i> var. <i>pelagica</i>	<i>H. rufescens</i>	12		~300	5	55	18	Correa-Reyes et al. 2009

Diatom	Abalone	Growth ($\mu\text{m d}^{-1}$)	Survival (%)	Initial SL (μm)	Initial Age (d)	Final Age (d)	Temp. ($^{\circ}\text{C}$)	Reference
<i>Nitzschia frustulum</i> var. <i>perminuta</i>	<i>H. rufescens</i>	16		~300	5	55	18	Correa-Reyes et al. 2009
<i>Nitzschia thermalis</i> var. <i>minor</i>	<i>H. rufescens</i>	29		~300	5	55	18	Correa-Reyes et al. 2009
Wild diatoms (<i>Navicula</i> sp. and <i>Cocconeis</i> sp. predominant) on macroalgae	<i>H. rufescens</i>	1.9%	29	<400	0	30	15	Muñoz et al. 2012
Wild diatoms (<i>Navicula</i> sp. and <i>Cocconeis</i> sp. predominant) on plastic plates	<i>H. rufescens</i>	0.8%	40	<400	0	30	15	Muñoz et al. 2012
<i>Amphora</i> sp.	<i>H. tuberculata</i> <i>coccinea</i>	50	~75	241	0	70	21	Courtois de Vicoise et al. 2012
<i>Navicula incerta</i>	<i>H. tuberculata</i> <i>coccinea</i>	46	~75	241	0	70	21	Courtois de Vicoise et al. 2012
<i>Nitzschia</i> sp.	<i>H. tuberculata</i> <i>coccinea</i>	43	~75	241	0	70	21	Courtois de Vicoise et al. 2012
<i>Proschkinia</i> sp.	<i>H. tuberculata</i> <i>coccinea</i>	38	~75	241	0	70	21	Courtois de Vicoise et al. 2012

Achnanthes brevipes was a remarkably poor diet and should be avoided for very young *H. kamtschatkana*. We saw no indication that postlarvae grazed on this diatom, and postlarvae died as quickly as those in the starvation control. At one point we observed an individual postlarva moving about its container but making no effort to feed on the *A. brevipes* cells over which it was moving. This was not due to cell size, as *A. brevipes* cells were not larger or smaller than other diatom diets tested. *Achnanthes brevipes* cells grow on short mucilage stalks (Toyoda et al. 2005), which might make them too difficult for *H. kamtschatkana* to consume, or the postlarvae may be responding to chemical cues indicating that the diatom is toxic or otherwise not suitable for consumption.

In comparison, *A. brevipes* may be a suitable feed for other, larger abalone species. *Haliotis diversicolor aquatilis* Reeve postlarvae under 1,000 μm shell length (Onitsuka et al. 2007) and *H. discus hannai* postlarvae less than 1,200 μm shell length (Takami et al. 2003) cannot detach the cells of related diatom *Achnanthes longipes* Agardh. At 4 weeks post-settlement with shell lengths of approximately 1,400 μm , however, *H. discus hannai* fed *A. longipes* and *A. brevipes* do grow in shell length (Kawamura et al. 1995). The greatest shell length of *H. kamtschatkana* in the current study was slightly less than 1,400 μm at 8.5 weeks, and no postlarvae fed *A. brevipes* survived to that point, let alone grew to that size.

Amphiprora paludosa, *N. incerta*, and *N. laevis* yielded moderate survival and growth. These three diatoms are all represented in previous literature but with variable performance as diets (Table 6). This may be related to cell density (Correa-Reyes et al. 2009). *Navicula incerta* is commonly a reference diet against which other diatom species are compared, or used as a fixed diet in tests of growing conditions for abalones, such as light intensity and seawater flow rate (Searcy-Bernal and Gorrostieta-Hurtado 2007; Correa-Reyes et al. 2009; Anguiano-Beltran

and Searcy-Bernal 2013). *Nitzschia laevis* was a more suitable food for *H. kamtschatkana* in our study than it is for *H. discus hannai* postlarvae, which cannot survive on it (Gordon et al. 2006).

Culture conditions and strain ID influence myriad diatom characteristics (e.g. Guerrini et al. 2000), and it is rare for any two studies to compare the same set of diatoms. Diatom adhesive strength is primarily measured in the context of testing anti-fouling coatings for underwater surfaces. These tests are focused on differences between coatings, not between diatoms, so their applicability to abalone postlarvae feeding constraints is limited. Likewise, studies of diatoms' organic composition often focus on only one or two diatoms at a time.

No previous work on yolk reserves in *H. kamtschatkana* exists. In our study, 15% of starved postlarvae survived to day 20, but only a single larva survived to day 26. Bacteria may have been present, but no diatoms or other algae. Even if bacteria were nutritionally available to postlarvae in the starvation control, without mucus secretions from diatoms, the bacteria would have limited growth media.

We used carbon-nitrogen ratio as a simple proxy for nutritional value of food. However, in this study we found no relationship between this ratio and postlarval survival or growth rates. In the snail *Potamopyrgus jenkinsi*, C:N ratios up to 15:1 were positively and tightly correlated with growth, but for higher ratios, the correlation was loose and negative (Dorgelo and Leonards 2001). In our study, only 2 diatoms had ratios of 15:1 or less and they did not follow this pattern: *C. closterium* (10.5:1 [SD, 3.7]) yielded higher growth than *A. paludosa* (9.7:1 [SD, 1.6]). For extensive consideration of the use of C:N ratios in molluscan mariculture, see Bayne (2009).

We recommend that future work on feeds for *H. kamtschatkana* postlarvae consider lipid and amino acid compositions. *Haliotis rufescens* postlarval growth rate correlates with amino acid

content but not fatty acid profiles (Correa-Reyes et al. 2009). In a study of 4 diatom diets, including *N. incerta*, *Amphora* sp. had the highest lipid and protein content, the lowest carbohydrate content, and yielded fastest growth of *H. tuberculata coccinea* Reeve postlarvae (Courtois de Vicose, Viera, et al. 2012). *Navicula incerta* yielded moderate growth, and showed moderate lipid, protein, and carbohydrate content; fatty acid profiles varied between all 4 diatoms in the study. Formulated feeds may be used to test different levels of biochemical components for abalone postlarval nutrition (Montaño-Vargas et al. 2005).

Unlike the other diatoms tested, *A. salina* was isolated from the middle of the geographic range of *H. kamtschatkana* (Saanich Inlet, British Columbia; NCMA 2019c), so we suggest that future workers test more diatoms isolated from this area. However, geography does not necessarily correlate with diet success, since *H. kamtschatkana* postlarvae grew well when fed *C. closterium* isolated from the Sargasso Sea in the North Atlantic open ocean (NCMA 2019a). *Navicula incerta* was isolated from San Francisco, near the southern end of *H. kamtschatkana*'s range (NCMA 2019d). The lowest performing diets in this experiment, *A. brevipes*, *A. paludosa*, and *N. laevis*, were all isolated from Nantucket Bay and Nantucket Sound, Massachusetts (NCMA 2019e; NCMA 2019b; NCMA 2019f).

***Haliotis kamtschatkana* growth rates**

Our study is the first quantitative account of growth rates of *H. kamtschatkana* postlarvae. Caldwell (1981) qualitatively described their growth as much slower than *H. rufescens*, which is a closely related species (Gallardo-Escarate et al. 2004; Crosson and Friedman 2018). In our study, mean shell length of postlarval *H. kamtschatkana* increased $11 \mu\text{m day}^{-1}$ on average (range 4.2 to $20 \mu\text{m day}^{-1}$) from days 7 to 61. In comparison, growth rate of postlarval *H.*

rufescens is approximately 10 to 30 $\mu\text{m day}^{-1}$ in the first 30 days post-settlement, and up to 125 $\mu\text{m day}^{-1}$ at day 60 post-settlement (Martinez-Ponce and Searcy-Bernal 1998; Gorrostieta-Hurtado and Searcy-Bernal 2004; Uriarte et al. 2006; Searcy-Bernal et al. 2007; Searcy-Bernal and Gorrostieta-Hurtado 2007; Correa-Reyes et al. 2009; Muñoz et al. 2012; Anguiano-Beltran and Searcy-Bernal 2013). *Haliotis kamtschatkana* postlarvae fed *C. closterium* in our study grew at 20 $\mu\text{m day}^{-1}$ once they were above $\sim 800 \mu\text{m}$ shell length; postlarvae of *H. discus hannai*, a species extensively cultivated in Asia and Chile, grow about 20 to 40 $\mu\text{m day}^{-1}$ when fed *C. closterium* at shell lengths of 800 to 1,200 μm (reviewed by Takami and Kawamura 2003), and postlarvae of *H. iris* are double the length of *H. kamtschatkana* at 60 days post-settlement when fed *C. closterium* (2,700 μm vs. 1,360 μm , respectively; Roberts, Kawamura, and Takami 1999). We also found that abalone growth rate increased exponentially with age over the course of our experiment. For example, from our model for *A. salina*, *N. incerta*, or *N. laevis*, growth rate was 6.2 $\mu\text{m day}^{-1}$ at day 7 and 16 $\mu\text{m day}^{-1}$ at day 61. Oddly, few other studies report growth as an exponential equation and instead give linear measures of growth, sometimes at different ages.

Shell length predicts soft tissue mass in gastropods generally (McKinney et al. 2004; Mehler et al. 2015) and in haliotids specifically (Najmudeen 2015). Strength of such relationships is variable both within and between species, and between size classes, but shell length is nonetheless very commonly used as a metric of abalone size. Shell length is suitable to our work because our goal is to support hatchery production of *H. kamtschatkana* up to a shell size suitable for restoration out-planting, in contrast to work increasing meat yields for abalones grown for food.

The radula

Radula development in *H. kamtschatkana* is similar to other species: the number of lateral teeth increases slowly; the number of marginal teeth increases rapidly; and shell length has a strong linear relationship with radula width, radula length, gap length between rows of teeth, and rachidian tooth size in postlarvae of *H. kamtschatkana* (Figure 7), *H. discus hannai* (Kawamura et al. 2001), *H. diversicolor aquatilis* (Onitsuka et al. 2004), and *H. iris* (Roberts, Kawamura, and Takami 1999). There were small differences between *H. kamtschatkana* and other species at the same shell lengths; for instance, the number of lateral teeth per row was slightly higher than it is in *H. discus hannai* (Kawamura et al. 2001), and the radula was narrower than it is in *H. iris* (Roberts, Kawamura, and Takami 1999). At shell lengths of 750 to 1,200 μm , the *H. kamtschatkana* radula was approximately one-quarter of the length of the shell, but in *H. iris* and *H. diversicolor aquatilis* of the same size it was approximately one-third of the length of the shell. (Roberts, Kawamura, and Takami 1999; Onitsuka et al. 2004).

Larval veliger age is strongly correlated with number of rows of teeth per radula (in *H. australis*, Moss 1999; in *H. discus hannai*, Takami et al. 2006), but rate of gain of tooth rows may (in *H. discus hannai*; Takami et al. 2006) or may not (in *Haliotis australis* Gmelin; Moss 1999) increase when a larva settles and metamorphoses. At day 10 post-fertilization, *H. australis* postlarvae consistently have 16 rows of teeth (Moss 1999). In our experiment, postlarvae at day 12 post-fertilization (day 5 post-settlement) had 17 rows of teeth, and, throughout our trial, grew approximately three-quarters as quickly as *H. australis* (Moss 1997).

Three measured characteristics did not increase with *H. kamtschatkana* shell length: buccal cartilage position, number of rows of teeth per radula, and the ratio between rachidian cusp length and rachidian height (Figure 7 and Table 5). Buccal cartilage position is associated with

the number of rows of teeth being used for feeding. Teeth posterior to the buccal cartilages are not in use but ready for deployment when anterior teeth wear out and are shed. Therefore, it appears that both number of teeth in use and teeth in standby are relatively constant at any given time in *H. kamtschatkana* postlarvae. This might indicate that rate of tooth wear is the same regardless of age or size in the first 60 days. The buccal cartilage position of other abalone species is unknown.

The ratio between rachidian cusp length and rachidian height is a proxy for rachidian tooth contact angle with the substratum. This ratio did not change with postlarval age or size (Figure 7, Table 5), nor did geometric morphometrics of landmark positions on teeth reveal any quantitative changes in tooth shape or relative position over time (Figure 8).

In general, as abalone age, individual teeth become gradually less serrated. However, we observed reduced serrations only in a 9-month old *H. kamtschatkana* juvenile outside of our feeding trial, not in postlarvae up to 61 days post-settlement, (Figure 7). Reduced tooth serrations are first visible at 1,200 μm shell length and 53 days post-settlement in *H. iris* (Roberts, Kawamura, and Takami 1999), 1,800 μm shell length in *H. diversicolor aquatilis* (Onitsuka et al. 2004), and 1,890 μm shell length and 49 days post-settlement in *H. discus hannai* (Kawamura et al. 2001). *Haliotis discus hannai* postlarvae at 17 days post-settlement and 1,145 μm shell length have teeth of similar size to our *H. kamtschatkana* postlarvae at 61 days post-settlement and similar shell length, but the former have teeth with a more delicate appearance and deeper serrations (Kawamura et al. 2001). If *H. kamtschatkana* tooth morphology mirrors that of other species, our study may have ended at around the time that decreased tooth serration would be noticeable.

Diatom shape, size, and adhesion did not seem to affect postlarval growth or survival in our study. The diatom with weakest adhesion, *A. paludosa*, resulted in unexceptional growth and survival. Film appearance and cell size were similar between *N. incerta*, *N. laevis*, and the near-deadly *A. brevipes*. We do not know the relative frustule strength of diatoms in our study, but *C. closterium* is known for a thin frustule (Reimann and Lewin 1964) and led to the fastest growth in our study. Our radula measurements did not explain success or failure of any of our diatom diets—radula morphology of *H. kamtschatkana* was comparable to other abalones that have succeeded when fed these same diets.

We found no evidence for morphological plasticity of the radula in response to diet. Plastic response to diet is known to occur in radulae of adult Littorinidae (Padilla 1998; Padilla 2004; Andrade and Solferini 2006; Molis et al. 2015) and other gastropods (e.g. *Placida dendritica*, Bleakney 1990; *Elysia viridis*, Jensen 1993); however, in some groups such plasticity does not occur (e.g. *Theodoxus fluviatilis*, Zettler et al. 2004; *Buccinanops globulosus*, Avaca et al. 2010; the polyplacophoran *Leptochiton asellus*, Sigwart and Carey 2014). Our abalone may have been too young to display plasticity, the study may have been too short, the diatoms too similar, or perhaps abalone in general do not have morphological plasticity of the radula.

Conclusion

We recommend that *H. kamtschatkana* hatcheries feed *A. salina* immediately after settlement, possibly paired with *C. closterium* to support growth. Future work should investigate diatom diets for *H. kamtschatkana* postlarvae during and immediately after settlement, and at 2 to 4 months post-settlement. This work should consider biochemical composition, and should reiterate diets with attention to whether it is the diatom species or the growing conditions that

drive diet success. *Amphora salina* may also be of interest to those studying settlement and metamorphosis success, because of the especially high early survival of postlarvae grown on it. Radula development in *H. kamtschatkana* is very similar to other abalone species, albeit slower, in tandem with overall growth. However, radula morphology did not illuminate why postlarval growth and survival were higher on some diatom diets than on others. *Haliotis kamtschatkana* postlarval success seemed driven mostly by abalone age and differences between diatom species, rather than any nuances of teeth.

LITERATURE CITED

- Adams DC, Collyer ML, Kaliontzopoulou A, Sherratt E. 2019. Geomorph: Software for geometric morphometric analyses. R package version 3.0.5. (R package). <https://cran.r-project.org/package=geomorph>.
- Allee WC. 1949. Principles of animal ecology. Philadelphia (PA): WB Saunders Company.
- Andrade SCS, Solferini VN. 2006. Transfer experiment suggests environmental effects on the radula of *Littoraria flava* (Gastropoda: Littorinidae). *J Molluscan Stud.* 72(1):111–116. doi:10.1093/mollus/eyi051.
- Anguiano-Beltran C, Searcy-Bernal R. 2013. Survival and growth of red abalone (*Haliotis rufescens*) postlarvae under different static and flow conditions. *J Shellfish Res.* 32(1):29–32. doi:10.2983/035.032.0106.
- Araya R, Bahamondes C, Barahona K, Silva-Aciaras F. 2010. Utilización de una biopelícula microalgal multiespecífica para optimizar la fijación larval y el crecimiento de abalón (*Haliotis rufescens*) en un criadero comercial. Application of multi-specific microalgae biofilm for optimization of the larvae settlement and growth of abalone (*Haliotis rufescens*) in a commercial hatchery. *Rev Biol Mar Oceanogr.* 45(1):59–69. doi:10.4067/S0718-19572010000100005.
- Argumedo-Hernández U, Siqueiros-Beltrones DA, Sánchez-Saavedra MDP. 2010. Nonselective diatom grazing by juveniles of red abalone (*Haliotis rufescens* Swainson, 1822) under experimental culture conditions. *J Shellfish Res.* 29(4):951–958. doi:10.2983/035.029.0429.
- Avaca MS, Martín PR, Narvarte MA. 2010. Description of the radula of *Buccinanops globulosus* (Neogastropoda, Nassariidae) and an assessment of its variability in northern Patagonian gulfs. *Malacologia.* 53(1):175–183. doi:10.4002/040.053.0111.
- Bates D, Mächler M, Bolker B, Walker S. 2015. Fitting linear mixed-effects models using lme4. *J Stat Softw.* 67(1):1–48. doi:10.18637/jss.v067.i01.
- Bayne BL. 2009. Carbon and nitrogen relationships in the feeding and growth of the Pacific oyster, *Crassostrea gigas* (Thunberg). *J Exp Mar Biol Ecol.* 374(1):19–30. doi:10.1016/j.jembe.2009.04.003.
- Bleakney J. 1990. Indirect evidence of a morphological response in the radula of *Placida dentritica* (Alder and Hancock, 1843) (Opisthobranchia: Ascoglossa/Sacoglossa) to different alga prey. *Veliger.* 33:111–115.
- Bouma JV. 2007. Early life history dynamics of pinto abalone (*Haliotis kamtschatkana*) and implications for recovery in the San Juan Archipelago, Washington State [Master of Science]. [Seattle (WA)]: University of Washington.

- Bouma JV, Rothaus DP, Straus KM, Vadopalas B, Friedman CS. 2012. Low juvenile pinto abalone *Haliotis kamtschatkana kamtschatkana* abundance in the San Juan Archipelago, Washington State. *Trans Am Fish Soc.* 141(1):76–83. doi:10.1080/00028487.2011.651551.
- de Brouwer JF., Stal LJ. 2002. Daily fluctuations of exopolymers in cultures of the benthic diatoms *Cylindrotheca closterium* and *Nitzschia* sp. (Bacillariophyceae). *J Phycol.* 38(3):464–472. doi:10.1046/j.1529-8817.2002.01164.x.
- Caldwell ME. 1981. Spawning, early development, and hybridization of *Haliotis kamtschatkana* Jonas [Master of Science]. [Seattle (WA)]: University of Washington.
- Campbell A, editor. 1999. Canadian Special Publication of Fisheries and Aquatic Sciences 130: Workshop on rebuilding abalone stocks in British Columbia. Ottawa (ON): NRC (National Research Council) Research Press.
<https://www.nrcresearchpress.com/doi/abs/10.1139/9780660182926>.
- Carson HS, Ulrich M. 2019. Status report for the pinto abalone in Washington. Olympia (WA): Washington Department of Fish and Wildlife. <https://wdfw.wa.gov/publications/02031>.
- Correa-Reyes JG, del Pilar Sanchez-Saavedra M, Teresa Viana M, Flores-Acevedo N, Vasquez-Pelaez C. 2009. Effect of eight benthic diatoms as feed on the growth of red abalone (*Haliotis rufescens*) postlarvae. *J Appl Phycol.* 21(4):387–393. doi:10.1007/s10811-008-9381-x.
- Courtois de Vicose G, Porta A, Viera MP, Fernandez-Palacios H, Izquierdo MS. 2012. Effects of density on growth rates of four benthic diatoms and variations in biochemical composition associated with growth phase. *J Appl Phycol.* 24(6):1427–1437. doi:10.1007/s10811-012-9799-z.
- Courtois de Vicose G, Viera MP, Huchette S, Izquierdo MS. 2012. Improving nursery performances of *Haliotis tuberculata coccinea*: nutritional value of four species of benthic diatoms and green macroalgae germlings. *Aquaculture.* 334:124–131. doi:10.1016/j.aquaculture.2011.12.040.
- Crosson LM, Friedman CS. 2018. Withering syndrome susceptibility of northeastern Pacific abalones: A complex relationship with phylogeny and thermal experience. *J Invertebr Pathol.* 151:91–101. doi:10.1016/j.jip.2017.11.005.
- Daume S, Krsinich A, Farrell S, Gervis M. 2000. Settlement, early growth and survival of *Haliotis rubra* in response to different algal species. *J Appl Phycol.* 12(3–5):479–488. doi:10.1023/A:1008110828581.
- Day R, Gilmour P, Huchette S. 2004. Effects of density and food supply on postlarval abalone: behaviour, growth and mortality. *J Shellfish Res.* 23(4):1009–1018.
- Ding JJ, Huang B, Hu YQ, Wang XB. 2017. The effects of different monospecific benthic diatoms on larval settlement, metamorphosis, survival, and growth of *Haliotis asinina* Linnaeus in the South China Sea. *Aquac Int.* 25(1):367–377. doi:10.1007/s10499-016-0035-8.

- Dorgelo J, Leonards PEG. 2001. Relationship between C/N ratio of food types and growth rate in the snail *Potamopyrgus jenkinsi* (E. A. Smith). *J North Am Benthol Soc.* 20(1):60–67. doi:10.2307/1468188.
- Ebert EE, Houk JL. 1984. Elements and innovations in the cultivation of red abalone *Haliotis rufescens*. *Aquaculture.* 39(1–4):375–392. doi:10.1016/0044-8486(84)90279-5.
- Fukazawa H, Takami H, Kawamura T, Watanabe Y. 2005. The effect of egg quality on larval period and postlarval survival of an abalone *Haliotis discus hannai*. *J Shellfish Res.* 24(4):1141–1147. doi:10.2983/0730-8000(2005)24[1141:TEOEQO]2.0.CO;2.
- Gallardo-Escarate C, Alvarez-Borrego J, Portilla MAD, Kober V. 2004. Karyotype of pacific red abalone *Haliotis rufescens* (Archaeogastropoda : Haliotidae) using image analysis. *J Shellfish Res.* 23(1):205–209.
- Geiger DL. 1999. Description of *Haliotis fatui* new species (Gastropoda : Vetigastropoda) from the tropical western Pacific. *Nautilus.* 113(3):73–77.
- Geiger DL. 2000. Distribution and biogeography of the Haliotidae (Gastropoda: Vetigastropoda) world-wide. *Boll Malacol.* 35:57–120.
- Gordon N, Neori A, Shpigel M, Lee J, Harpaz S. 2006. Effect of diatom diets on growth and survival of the abalone *Haliotis discus hannai* postlarvae. *Aquaculture.* 252(2–4):225–233. doi:10.1016/j.aquaculture.2005.06.034.
- Gorrostieta-Hurtado E, Searcy-Bernal R. 2004. Combined effects of light condition (constant illumination or darkness) and diatom density on postlarval survival and growth of the abalone *Haliotis rufescens*. *J Shellfish Res.* 23(4):1001–1008.
- Guerrini F, Cangini M, Boni L, Trost P, Pistocchi R. 2000. Metabolic responses of the diatom *Achnanthes brevipes* (Bacillariophyceae) to nutrient limitation. *J Phycol.* 36(5):882–890. doi:10.1046/j.1529-8817.2000.99070.x.
- Hickman C. 1981. Evolution and function of asymmetry in the archeogastropod radula. *Veliger.* 23(3):189–194.
- Hoagland KD, Rosowski JR, Gretz MR, Roemer SC. 1993. Diatom extracellular polymeric substances: function, fine structure, chemistry, and physiology. *J Phycol.* 29:537–566. doi:10.1111/j.0022-3646.1993.00537.x.
- Holland R, Dugdale TM, Wetherbee R, Brennan AB, Finlay JA, Callow JA, Callow ME. 2004. Adhesion and motility of fouling diatoms on a silicone elastomer. *Biofouling.* 20(6):323–329. doi:10.1080/08927010400029031.
- Jensen KR. 1993. Morphological adaptations and plasticity of radular teeth of the Sacoglossa (= Ascoglossa) (Mollusca: Opisthobranchia) in relation to their food plants. *Biol J Linn Soc.* 48(2):135–155. doi:10.1111/j.1095-8312.1993.tb00883.x.

- Johnston D, Moltschaniwskyj N, Wells J. 2005. Development of the radula and digestive system of juvenile blacklip abalone (*Haliotis rubra*): potential factors responsible for variable weaning success on artificial diets. *Aquaculture*. 250(1–2):341–355. doi:10.1016/j.aquaculture.2005.03.012.
- Joyce A, Utting S. 2015. The role of exopolymers in hatcheries: an overlooked factor in hatchery hygiene and feed quality. *Aquaculture*. 446:122–131. doi:10.1016/j.aquaculture.2015.04.037.
- Kawamura T, Hirano R. 1992. Seasonal changes in benthic diatom communities colonizing glass slides in Aburatsubo Bay, Japan. *Diatom Res*. 7(2):227–239. doi:10.1080/0269249X.1992.9705216.
- Kawamura T, Roberts RD, Nicholson CM. 1998. Factors affecting the food value of diatom strains for post-larval abalone *Haliotis iris*. *Aquaculture*. 160(1–2):81–88. doi:10.1016/S0044-8486(97)00223-8.
- Kawamura T, Roberts RD, Takami H. 1998. A review of the feeding and growth of postlarval abalone. *J Shellfish Res*. 17(3):615–625.
- Kawamura T, Saïdo T, Takami H, Yamashita Y. 1995. Dietary value of benthic diatoms for the growth of post-larval abalone *Haliotis discus hannai*. *J Exp Mar Biol Ecol*. 194(2):189–199. doi:10.1016/0022-0981(95)00099-2.
- Kawamura T, Takami H. 1995. Analysis of feeding and growth rate of newly metamorphosed abalone *Haliotis discus hannai* fed on 4 species of benthic diatom. *Fish Sci*. 61(2):357–358. doi:10.2331/fishsci.61.357.
- Kawamura T, Takami H, Roberts RD, Yamashita Y. 2001. Radula development in abalone *Haliotis discus hannai* from larva to adult in relation to feeding transitions. *Fish Sci*. 67(4):596–605. doi:10.1046/j.1444-2906.2001.00295.x.
- Kuznetsova A, Brockhoff PB, Christensen RHB. 2017. lmerTest package: Tests in linear mixed effects models. *J Stat Softw*. 82(13):1–26. doi:10.18637/jss.v082.i13.
- Martinez-Ponce DR, Searcy-Bernal R. 1998. Grazing rates of red abalone (*Haliotis rufescens*) postlarvae feeding on the benthic diatom *Navicula incerta*. *J Shellfish Res*. 17(3):627–630.
- McKinney RA, Glatt SM, Williams SR. 2004. Allometric length-weight relationships for benthic prey of aquatic wildlife in coastal marine habitats. *Wildl Biol*. 10(1):241–249. doi:10.2981/wlb.2004.029.
- Mehler K, Acharya K, Sada DW. 2015. Spatial and Temporal Pattern in Length-Mass Regressions of Freshwater Gastropods in Nevada Spring Ecosystems. *Malacologia*. 58(1–2):167–177. doi:10.4002/040.058.0205.

- Molis M, Scrosati RA, El-Belely EF, Lesniowski TJ, Wahl M. 2015. Wave-induced changes in seaweed toughness entail plastic modifications in snail traits maintaining consumption efficacy. *J Ecol.* 103(4):851–859. doi:10.1111/1365-2745.12386.
- Montaño-Vargas J, Viana MT, D’Abramo LR, Shimada A, Vásquez-Pelaez C. 2005. Growth and energy utilization of juvenile pink abalone *Haliotis corrugata* fed diets containing different levels of protein and two starch:lipid ratios. *J Shellfish Res.* 24(4):1179–1185.
- Moss GA. 1997. Early juvenile growth of the abalone *Haliotis australis* in culture. *Molluscan Res.* 18(2):153–159. doi:10.1080/13235818.1997.10673689.
- Moss GA. 1999. Factors affecting settlement and early post-settlement survival of the New Zealand abalone *Haliotis australis*. *N Z J Mar Freshw Res.* 33(2):271–278. doi:10.1080/00288330.1999.9516876.
- Muñoz P, Ambler R, Bulboa C. 2012. Settlement, survival, and post-larval growth of red abalone, *Haliotis rufescens*, on polycarbonate plates treated with germlings of *Ulva* sp. *J World Aquac Soc.* 43(6):890–895. doi:10.1111/j.1749-7345.2012.00615.x.
- Najmudeen TM. 2015. Biometric relationships of the Indian abalone *Haliotis varia* Linnaeus 1758 from Mandapam waters of Gulf of Mannar, south-east coast of India. *Indian J Fish.* 62:146–150.
- Najmudeen TM. 2017. Competitive interactions between two marine benthic microalgae in mixed culture. *Indian J Geo-Mar Sci.* 46(7):1417–1422.
- National Center for Marine Algae and Microbiota. 2019a. CCMP340 | NCMA - Culturing Diversity. Bigelow Natl Cent Mar Algae Microbiota. [accessed 2018 Feb 23]. <https://ncma.bigelow.org/ccmp340>.
- National Center for Marine Algae and Microbiota. 2019b. CCMP559 | NCMA - Culturing Diversity. Bigelow Natl Cent Mar Algae Microbiota. [accessed 2018 Feb 23]. <https://ncma.bigelow.org/ccmp559>.
- National Center for Marine Algae and Microbiota. 2019c. CCMP1119 | NCMA - Culturing Diversity. Bigelow Natl Cent Mar Algae Microbiota. [accessed 2018 Feb 23]. <https://ncma.bigelow.org/ccmp1119>.
- National Center for Marine Algae and Microbiota. 2019d. CCMP542 | NCMA - Culturing Diversity. Bigelow Natl Cent Mar Algae Microbiota. [accessed 2018 Feb 23]. <https://ncma.bigelow.org/ccmp542>.
- National Center for Marine Algae and Microbiota. 2019e. CCMP100 | NCMA - Culturing Diversity. Bigelow Natl Cent Mar Algae Microbiota. [accessed 2018 Feb 23]. <https://ncma.bigelow.org/ccmp100>.

National Center for Marine Algae and Microbiota. 2019f. CCMP125 | NCMA - Culturing Diversity. Bigelow Natl Cent Mar Algae Microbiota. [accessed 2018 Feb 23]. <https://ncma.bigelow.org/ccmp125>.

Neuman MJ, Wang S, Busch S, Friedman C, Gruenthal K, Gustafson R, Kushner D, Stierhoff K, Vanblaricom G, Wright S. 2018. A status review of pinto abalone (*Haliotis kamtschatkana*) along the west coast of North America: Interpreting trends, addressing uncertainty, and assessing risk for a wide-ranging marine invertebrate. *J Shellfish Res.* 37(4):869–910. doi:10.2983/035.037.0415.

Onitsuka T, Kawamura T, Ohashi S, Horii T, Watanabe Y. 2004. Morphological changes in the radula of abalone *Haliotis diversicolor aquatilis* from post-larva to adult. *J Shellfish Res.* 23(4):1079–1085.

Onitsuka T, Kawamura T, Ohashi S, Horii T, Watanabe Y. 2007. Dietary value of benthic diatoms for post-larval abalone *Haliotis diversicolor* associated with feeding transitions. *Fish Sci.* 73(2):295–302. doi:10.1111/j.1444-2906.2007.01335.x.

Padilla DK. 1998. Inducible phenotypic plasticity of the radula in *Lacuna* (Gastropoda: Littorinidae). *Veliger.* 41(2):201–204.

Padilla DK. 2004. Form and function of radular teeth of herbivorous molluscs: focus on the future. *Am Malacol Bull.* 18(1–2):163–168.

Pang SJ, Zhang ZH, Bao Y, Gao SQ. 2006. Settling abalone veliger larvae in a free-swimming microalgal culture. *Aquaculture.* 258(1–4):327–336. doi:10.1016/j.aquaculture.2006.03.028.

Paul AJ, Paul JM. 1981. Temperature and growth of maturing *Haliotis kamtschatkana* Jonas. *Veliger.* 23(4):321–324.

Paul AJ, Paul JM. 1998. Respiration rate and thermal tolerances of pinto abalone *Haliotis kamtschatkana*. *J Shellfish Res.* 17(3):743–745.

Ponder WF, Lindberg DR. 1997. Towards a phylogeny of gastropod molluscs: an analysis using morphological characters. *Zool J Linn Soc.* 119(2):83–265. doi:10.1111/j.1096-3642.1997.tb00137.x.

Rasband W. 1997. ImageJ. Bethesda (MD): National Institutes of Health (US). <http://imagej.nih.gov/ij/>.

Rasmussen K, Østgaard K. 2001. Adhesion of the marine fouling diatom *Amphora coffeaeformis* to non-solid gel surfaces. *Biofouling.* 17(2):103–115. doi:10.1080/08927010109378470.

Ravizza M, Hallegraeff G. 2015. Environmental conditions influencing growth rate and stalk formation in the estuarine diatom *Licmophora flabellata* (Carmichael ex Greville) C. Agardh. *Diatom Res.* 30(2):197–208. doi:10.1080/0269249X.2015.1020071.

Reimann BEF, Lewin JC. 1964. The diatom genus *Cylindrotheca* Rabenhorst. J R Microsc Soc. 83(3):283–296. doi:10.1111/j.1365-2818.1964.tb00542.x.

Roberts RD, Kawamura T, Nicholson CM. 1999. Growth and survival of postlarval abalone (*Haliotis iris*) in relation to development and diatom diet. J Shellfish Res. 18(1):243–250.

Roberts RD, Kawamura T, Takami H. 1999. Morphological changes in the radula of abalone (*Haliotis iris*) during post-larval development. J Shellfish Res. 18(2):637–644.

Roberts RD, Lapworth C, Barker RJ. 2001. Effect of starvation on the growth and survival of post-larval abalone (*Haliotis iris*). Aquaculture. 200(3–4):323–338. doi:10.1016/S0044-8486(01)00531-2.

Rothaus DP, Vadopalas B, Friedman CS. 2008. Precipitous declines in pinto abalone (*Haliotis kamtschatkana kamtschatkana*) abundance in the San Juan Archipelago, Washington, USA, despite statewide fishery closure. Can J Fish Aquat Sci. 65(12):2703–2711. doi:10.1139/F08-168.

Searcy-Bernal R, Gorrostieta-Hurtado E. 2007. Effect of darkness and water flow rate on survival, grazing and growth rates of abalone *Haliotis rufescens* postlarvae. J Shellfish Res. 26(3):789–794. doi:10.2983/0730-8000(2007)26[789:EODAWF]2.0.CO;2.

Searcy-Bernal R, Pérez-Sánchez E, Anguiano-Beltrán C, Flores-Aguilar R. 2007. Metamorphosis and postlarval growth of abalone *Haliotis* [sic] *rufescens* in a Mexican commercial hatchery. J Shellfish Res. 26(3):783–787. doi:10.2983/0730-8000(2007)26[783:MAPGOA]2.0.CO;2.

Searcy-Bernal R, Velez-Espino LA, Anguiano-Beltran C. 2001. Effect of biofilm density on grazing and growth rates of *Haliotis fulgens* postlarvae. J Shellfish Res. 20(2):587–591.

Sigwart JD, Carey N. 2014. Grazing under experimental hypercapnia and elevated temperature does not affect the radula of a chiton (Mollusca, Polyplacophora, Lepidopleurida). Mar Environ Res. 102:73–77. doi:10.1016/j.marenvres.2014.05.004.

Steneck RS, Watling L. 1982. Feeding capabilities and limitation of herbivorous molluscs: a functional group approach. Mar Biol. 68(3):299–319. doi:10.1007/BF00409596.

Takami H, Kawamura T. 2003. Dietary changes in the abalone, *Haliotis discus hannai*, and relationship with the development of the digestive organ. Jpn Agric Res Q JARQ. 37(2):89–98. doi:10.6090/jarq.37.89.

Takami H, Kawamura T, Yamashita Y. 1997. Survival and growth rates of post-larval abalone *Haliotis discus hannai* fed conspecific trail mucus and/or benthic diatom *Cocconeis scutellum* var. *parva*. Aquaculture. 152(1–4):129–138. doi:10.1016/S0044-8486(96)01524-4.

Takami H, Kawamura T, Yamashita Y. 1998. Development of polysaccharide degradation activity in postlarval abalone *Haliotis discus hannai*. J Shellfish Res. 17:723–727.

- Takami H, Kawamura T, Yamashita Y. 2000. Starvation tolerance of newly metamorphosed abalone *Haliotis discus hannai*. Fish Sci. 66:1180–1182. doi:10.1046/j.1444-2906.2000.00186.x.
- Takami H, Muraoka D, Kawamura T, Yamashita Y. 2003. When is the abalone *Haliotis discus hannai* Ino 1953 first able to use brown macroalgae? J Shellfish Res. 22(3):795–800.
- Takami H, Oshino A, Sasaki R, Fukazawa H, Kawamura T. 2006. Age determination and estimation of larval period in field caught abalone (*Haliotis discus hannai* Ino 1953) larvae and newly metamorphosed post-larvae by counts of radular teeth rows. J Exp Mar Biol Ecol. 328(2):289–301. doi:10.1016/j.jembe.2005.07.014.
- Tesson B, Hildebrand M. 2013. Characterization and localization of insoluble organic matrices associated with diatom cell walls: Insight into their roles during cell wall formation. PLOS ONE. 8(4):e61675. doi:10.1371/journal.pone.0061675.
- Toyoda K, Cox EJ, Sims PA, Williams DM. 2005. The typification of *Achnanthes bory* based on *Echinella stipitata* Lyngbye, with an account of the morphology and fine structure of Lyngbye's species. Diatom Res. 20(2):375–386. doi:10.1080/0269249X.2005.9705643.
- Uriarte I, Roberts R, Farias A. 2006. The effect of nitrate supplementation on the biochemical composition of benthic diatoms and the growth and survival of post-larval abalone. Aquaculture. 261(1):423–429. doi:10.1016/j.aquaculture.2006.08.007.
- Vadopalas B, Watson J. 2014. Recovery plan for pinto abalone (*Haliotis kamtschatkana*) in Washington state (using a collaborative management approach). Bainbridge Island, WA: Puget Sound Restoration Fund.
http://www.restorationfund.org/sites/default/files/2014_Kamtschatkana_Recovery_Plan.pdf.
- Viana MT, Correa G, Lazo JP, Frías-Díaz R, Durazo-Beltrán E, Vasquez-Pelaez C. 2007. Digestive physiology and metabolism of green abalone *Haliotis fulgens* from postlarvae to juvenile, fed three different diatoms. Aquaculture. 271(1–4):449–460. doi:10.1016/j.aquaculture.2007.04.072.
- Wallace SS. 1999. Evaluating the effects of three forms of marine reserve on northern abalone populations in British Columbia, Canada. Conserv Biol. 13(4):882–887. doi:10.1046/j.1523-1739.1999.98117.x.
- Wang P, Park BS, Kim JH, Kim J-H, Lee H-O, Han M-S. 2014. Phylogenetic position of eight *Amphora sensu lato* (Bacillariophyceae) species and comparative analysis of morphological characteristics. Algae. 29(2):57–73. doi:10.4490/algae.2014.29.2.057.
- Xing R, Wang C, Cao X, Chang Y. 2007. Settlement, growth and survival of abalone, *Haliotis discus hannai*, in response to eight monospecific benthic diatoms. J Appl Phycol. 20(1):47–53. doi:10.1007/s10811-007-9179-2.

Xu W, Wang J, Tan L, Guo X, Xue Q. 2019. Variation in allelopathy of extracellular compounds produced by *Cylindrotheca closterium* against the harmful-algal-bloom dinoflagellate *Prorocentrum donghaiense*. *Mar Environ Res.* 148:19–25. doi:10.1016/j.marenvres.2019.05.005.

Zettler ML, Frankowski J, Bochert R, Rohner M. 2004. Morphological and ecological features of *Theodoxus fluviatilis* (Linnaeus, 1758) from Baltic brackish water and German freshwater populations. *J Conchol.* 38:305–316.

Zhang Y, Gao Y, Liang J, Chen C, Zhao D, Li X, Li Y, Wu W. 2010. Diatom diet selectivity by early post-larval abalone *Haliotis diversicolor supertexta* under hatchery conditions. *Chin J Oceanol Limnol.* 28(6):1187–1194. doi:10.1007/s00343-010-0019-x.